# **Facts & Myths About Cataract**



#### **MYTH: Only older Americans develop cataract**

**FACT:** While cataract affects more than 24 million Americans age 40 and older, cataract can occur among young adults or children. Risk factors that may lead to getting cataract at a younger age include:

- Intense heat or long-term exposure to UV rays from the sun
- Certain diseases, such as diabetes
- Inflammation in the eye
- Hereditary influences
- Events before birth, such as German measles in the mother
- Long-term steroid use
- Severe long-term nearsightedness (myopia)
- Eye injuries
- Eye diseases
- Smoking

#### **MYTH: Taking Vitamin E or Vitamin C can** prevent cataract

**FACT:** Some research centers are studying the link between these vitamins and cataract prevention. However, it will be many years before the studies can determine if vitamin C or E actually reduces a person's risk of cataract. Until then, it's best not to take these vitamins in large doses unless they have been prescribed by your physician.

# MYTH: The best time to have cataract surgery done is when it is first diagnosed.

**FACT:** Cataract removal is elective surgery, which means it is the patient's choice when to undergo the procedure. Most people need surgery when the cataract causes enough vision loss to interfere with work, play or other day-to-day tasks. You, your eye doctor, and family members should decide together when and if surgery is needed.



over 🖌

This publication is copyrighted. This sheet may be reproduced—unaltered in hard print (photocopied) for educational purposes only. The Prevent Blindness America name, logo, telephone number and copyright information may not be omitted. Electronic reproduction, other reprint, excerption or use is not permitted without written consent. Because of the time-sensitive nature of the information contained in this publication, contact Prevent Blindness America for updates.

### **MYTH: Taking aspirin can prevent cataract**

**FACT:** There are not enough facts or evidence to say whether aspirin prevents cataract. Aspirin in large doses can be harmful. Unless your doctor prescribes aspirin for an ailment, it's best to avoid taking this medication on a regular basis.

#### **MYTH: Lasers are used to remove cataract**

**FACT:** In cataract treatment, the clouded lens is surgically removed and then replaced with an artificial lens implant. If a patient has cataract in both eyes, separate surgeries are scheduled. Sometimes the membrane behind the implant may become cloudy after cataract surgery. Laser treatment then may be used to open up the cloudy membrane.

## **MYTH: Cataract can be treated with eye drops**

**FACT:** Surgery is the only proven treatment for cataract. Cataract cannot be treated with medicines.

#### **MYTH: Cataract surgery is dangerous**

**FACT:** Cataract surgery is a delicate operation. Yet, it is one of the safest operations done today. More than 95% of surgeries are successful. Fewer than 5% of cases have complications such as inflammation, bleeding, infection and retinal detachment.

#### MYTH: It can take months to recover from cataract surgery

**FACT:** In many cases, patients often can see well enough to resume normal activities a few days after having cataract surgery. Your vision will continue to improve over the following weeks and months. However, if you have additional eye problems, such as glaucoma, your recovery time might take longer.

